Drug Information Sheet("Kusuri-no-Shiori")

External
Revised: 09/2026

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:Latanoprost Ophthalmic Solution 0.005% "WAKAMOTO"
Active ingredient: Latanoprost
Dosage form:limpid and colorless ophthalmic solution, 2.5mL, (cap) pearl white
Imprint or print on wrapping:



Effects of this medicine

This medicine lowers eye pressure by increasing fluid drainage from the eye.
It is usually used to treat glaucoma and ocular hypertension.

The following patients may need to be careful when using this medicine.Be sure to tell your doctor and pharmacist.

・If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
If you have: an eye with aphakia or intraocular lens, bronchial asthma/history of bronchial asthma, intraocular inflammation (iritis or uveitis) , possible herpesvirus latency or angle-closure glaucoma.
If you wear contact lenses.
・If you are pregnant or breastfeeding.
・If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

・Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
・In general, apply 1 drop at a time, once a day. Strictly follow the instructions.
・Make sure not to let the tip of the container touch any part of your eye.
・After applying, keep your eyes closed without blinking for a while (1 to 5 minutes).
・If this medicine gets on your eyelid, wipe it off immediately.
・If you need to apply other eye drops at the same time, wait more than 5 minutes.
・If you are wearing contact lenses, take off the lenses and then administer. Wait more than 15 minutes to wear your contact lenses again. (It might lead contact lenses to discoloration.)
・If you miss a dose, apply 1 drop as soon as possible in case of the same day. If you notice on the next day, skip the missed dose and continue your regular dosing schedule. You should not apply twice a day or 2 drops at one time.
・If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
・Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

・Applying the eye-drops may cause iris pigmentation, followed by gradual increase of the symptom. The symptom may stop after withdrawal of the eye-drops, but there may be difference in color tone compared to the other eye without eye-drop treatment. If you become aware of it, consult with your doctor.
・This medicine may cause transient blurred vision. Avoid driving a car or operating machinery until these symptoms recover.

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include conjunctival hyperaemia, ocular irritancy symptom, itch, eye pain, blepharitis (eye lid infection), eyelid pigmentation, foggy vision, abnormal eyelash, reduced visual acuity, lacrimation, headache, throat discomfort and nausea. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

・pupil color change [iris pigmentation]

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

・The amount of contents seems to be insufficient because 2.5mL of solution is packed in a 5mL plastic bottle.
・Keep out of the reach of children. Store in the refrigerator (2 to 8 degrees Celsius) away from light.
・Discard the remainder that has passed 1 month after opening without storing it. If you do not know how to dispose of it, please consult the pharmacy or medical institution where you received it. Do not give it to others.

For healthcare professional use only / /

For further information, talk to your doctor or pharmacist.