Drug Information Sheet("Kusuri-no-Shiori")

Injection Published: 06/2019

	Fublished. 00/2019
The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.	
Brand name:LIMOLOSAL INJECTION 20mg(0.4%)	in .
Active ingredient: Betamethasone sodium phosphate	
Dosage form: Injection	A
Print on wrapping:	A Designed of the second se
	LAKEN -
Effects of this medicine	
This medicine is a synthetic corticosteroid preparation, and it shows anti-inflammatory effe	
immunosuppressive effects and various metabolic effects. It is usually used used for the treat	ment of various diseases
in each department such as internal medicine, pediatrics and surgery.	
However, it does not cure the causes of these diseases in themselves.	
Before using this medicine, be sure to tell your doctor and pharmacist	
•If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicine	S.
If you have infection or a joint with abnormal mobility.	
If you have generalized mycosis, peptic ulcer, psychosis, tuberculosis, herpes simplex keratitis, posterior	
subcapsular cataract, glaucoma, hypertension, abnormal electrolyte or thrombosis.	
If you have recently had a surgery in visceral organ.	
If you have a history of acute myocardial infarction.	
If you have viral conjunctiva and corneal disease, tuberculosis eye disease, fungal eye disea	se and acute suppurative
eye disease.	
• If you are pregnant or breastfeeding.	
• If you are taking any other medicinal products. (Some medicines may interact to enhance of	or diminish medicinal
effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)	
Dosing schedule (How to take this medicine)	prescription medicines.
• Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))	
•In general, this medicine is injected or infused intravenously, by intravenous drip, intramus	
cavity, into spinal cord cavity, into soft tissues, into tendon sheaths, intranasally etc., or it is administered by	
using a nebulizer. The usage varies according to your diseases.	it is administered by
•The treatment span is determined according to your symptoms. Consult with your doctor about your schedule for	
use.	about your schedule for
Precautions while taking this medicine	
• Pay attention to not to be infected with chickenpox or measles. Consult with your doctor	immediately if you are
suspected to have these infections, consult with your doctor immediately.	miniculatory in you are
Subpoolou to have these inconsistent with your usetor ininiouratory.	
Possible adverse reactions to this medicine	
The most commonly reported adverse reactions include malaise, loss of appetite, nausea, mo	on face and instability of
the joint (when intraarticular injection). If any of these symptoms occur, consult with your doctor or pharmacist.	
The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated	
in brackets. If any of these symptoms occur, stop taking this medicine and see yo	
immediately.	
•cold-like symptoms, malaise, fever [induced infection, exacerbation of infection]	
•nausea, black stool, abdominal pain [gastrointestinal tract ulcer, gastrointestinal perforation of the store	on, pancreatitis]
•feeling depressed, feeling unmotivated, spasmodic muscular contraction [psychiatric distur	
convulsion]	
•low back pain and/or back pain, joint pain, muscle stiffness [osteoporosis, aseptic necrosis	s of bone head in femur.
humerus, myopathy]	,
The above symptoms do not describe all the adverse reactions to this medicine. Consult with your	
doctor or pharmacist if you notice any symptoms of concern other than those listed above.	
Storage conditions and other information	
•When you recieve a live vaccine, be sure to consult with your doctor.	
For healthcare professional use only / /	

For further information, talk to your doctor or pharmacist.