

Drug Information Sheet("Kusuri-no-Shiori")

External

Published: 12/2025

The information on this sheet is based on approvals granted by the Japanese regulatory authority. Approval details may vary by country. Medicines have adverse reactions (risks) as well as efficacies (benefits). It is important to minimize adverse reactions and maximize efficacy. To obtain a better therapeutic response, patients should understand their medication and cooperate with the treatment.

Brand name:EPINASTINE HYDROCHLORIDE LX OPHTHALMIC SOLUTION 0.1% "WAKAMOTO"



Active ingredient:Epinastine hydrochloride

Dosage form:limpid and colorless ophthalmic solution, 5mL, (cap) dark orange

Imprint or print on wrapping:

Effects of this medicine

This medicine suppresses allergic symptoms such as itchy/bloodshot eyes by histamine H₁-receptor antagonism and suppressing release of allergy-causing substances such as histamine.

It is usually used to treat allergic conjunctivitis.

The following patients may need to be careful when using this medicine. Be sure to tell your doctor and pharmacist.

- If you have previously experienced any allergic reactions (itch, rash, etc.) to any medicines or foods.
- If you wear contact lenses.
- If you are pregnant or breastfeeding.
- If you are taking any other medicinal products. (Some medicines may interact to enhance or diminish medicinal effects. Beware of over-the-counter medicines and dietary supplements as well as other prescription medicines.)

Dosing schedule (How to take this medicine)

- Your dosing schedule prescribed by your doctor is((to be written by a healthcare professional))
- In general, apply 1 drop at a time, 2 times a day (morning and evening). Strictly follow the instructions.
- Make sure not to let the tip of the container touch any part of your eye.
- After applying, keep your eyes closed without blinking for a while (1 to 5 minutes).
- If this medicine gets on your eyelid, wipe it off immediately.
- If you need to apply other eye drops at the same time, wait more than 5 minutes.
- If you are wearing soft contact lenses, take off the lenses and then apply. After an interval of more than 5 minutes to wear contact lenses again.
- If you miss a dose, apply the missed dose as soon as possible. However, if it is almost time for the next dose, skip the missed dose and follow your regular dosing schedule. You should not apply two doses at one time.
- If you accidentally apply more than your prescribed dose, consult with your doctor or pharmacist.
- Do not stop applying this medicine unless your doctor instructs you to do so.

Precautions while taking this medicine

Possible adverse reactions to this medicine

The most commonly reported adverse reactions include bloodshot eyes. If any of these symptoms occur, consult with your doctor or pharmacist.

The symptoms described below are rarely seen as initial symptoms of the adverse reactions indicated in brackets. If any of these symptoms occur, stop taking this medicine and see your doctor immediately.

No pertinent entries.

The above symptoms do not describe all the adverse reactions to this medicine. Consult with your doctor or pharmacist if you notice any symptoms of concern other than those listed above.

Storage conditions and other information

- Keep out of reach of children. Store at room temperature (1 to 30 degrees Celsius) away from direct sunlight and moisture.
- Discard the remainder that has passed 1 month after opening without storing it. If you do not know how to dispose of it, consult the pharmacy or medical institution where you received it. Do not give it to others.

For healthcare professional use only

/ /

For further information, talk to your doctor or pharmacist.